Serial No.: 09/830,144 Filed: April 20, 2001

Page : 3 of 14

AMENDMENTS TO THE CLAIMS

Claims 2-67 are pending.

Claims 2-67 are being canceled without prejudice or disclaimer, and new claims 68-87 are being added.

After the amendments, claims 68-87 will be pending.

This listing of claims replaces all prior versions and listings of claims in the application:

<u>Listing of Claims</u>:

1.-67. (Canceled)

- 68. (New) A method for screening compounds inhibiting production of an inflammatory cytokine, comprising:
- (a) providing a sample comprising a TAK1 and a TAB1;
- (b) contacting a test compound with the TAK1 and the TAB1;
- (c) detecting binding between the TAK1 and the TAB1; and
- (d) selecting a compound that inhibits the binding; wherein the TAK1 of (a) is selected from the group consisting of
 - (i) a protein comprising amino acids 76 to 303 of SEQ ID NO:2; and
- (ii) a protein that binds to amino acids 437 to 504 of SEQ ID NO:4 and comprises an amino acid sequence encoded by a DNA sequence that hybridizes with the complement of nucleotides 408 to 1091 of SEQ ID NO:1 under washing conditions of 42°C, 5 x SSC, 0.1% sodium dodecyl sulfate, and 50% formamide;

and wherein the TAB1 of (a) is selected from the group consisting of

- (iii) a protein comprising amino acids 437 to 504 of SEQ ID NO:4; and
- (iv) a protein that binds to amino acids 76 to 303 of SEQ ID NO:2 and comprises an amino acid sequence encoded by a DNA that hybridizes with the complement of

Serial No.: 09/830,144 Filed: April 20, 2001

Page : 4 of 14

nucleotides 1338 to 1541 of SEQ ID NO:3 under washing conditions of 42°C, 5 x SSC, 0.1% sodium dodecyl sulfate, and 50% formamide.

- 69. (New) The method of claim 68, wherein the TAK1 and/or the TAB1 is fused with a peptide.
- 70. (New) The method of claim 68, wherein the TAK1 or the TAB1 is linked to a support.
- 71. (New) The method of claim 68, wherein a label is attached to the TAK1 or the TAB1 and the binding is detected by detecting or measuring the label.
- 72. (New) The method of claim 68, wherein the binding is detected by detecting or measuring the TAB1 bound to the TAK1 with a primary antibody against TAB1 or a primary antibody against a peptide fused with the TAB1.
- 73. (New) The method of claim 68, wherein the binding is detected by detecting or measuring the TAK1 bound to the TAB1 with a primary antibody against TAK1 or a primary antibody against a peptide fused with the TAK1.
- 74. (New) The method of claim 68, wherein the binding is detected by detecting or measuring the TAB1 bound to the TAK1 with a primary antibody against the TAB1 or a primary antibody against a peptide fused with TAB1, and a secondary antibody against the primary antibody.
- 75. (New) The method of claim 68, wherein the binding is detected by detecting or measuring the TAK1 bound to the TAB1 with a primary antibody against TAK1 or a primary antibody against a peptide fused with the TAK1, and a secondary antibody against the primary antibody.

Serial No.: 09/830,144 Filed: April 20, 2001

Page : 5 of 14

76. (New) The method of claim 74, wherein the primary antibody or the secondary antibody is labeled with a radioisotope, enzyme, or fluorescent substance.

- 77. (New) The method of claim 68, wherein the TAK1 of (a) comprises amino acids 76 to 303 of SEQ ID NO:2.
- 78. (New) The method of claim 68, wherein the TAK1 of (a) is a protein that binds to amino acids 437 to 504 of SEQ ID NO:4 and comprises an amino acid sequence encoded by a DNA that hybridizes with the complement of nucleotides 408 to 1091 of SEQ ID NO:1 under washing conditions of 42°C, 5 x SSC, 0.1% sodium dodecyl sulfate, and 50% formamide.
- 79. (New) The method of claim 68, wherein the TAK1 of (a) is a protein that binds to amino acids 437 to 504 of SEQ ID NO:4 and comprises an amino acid sequence that is encoded by a DNA that hybridizes with the complement of nucleotides 408 to 1091 of SEQ ID NO:1 under washing conditions of 60°C, 0.1 x SSC, and 0.1% sodium dodecyl sulfate.
- 80. (New) The method of claim 68, wherein the TAB1 of (a) comprises amino acids 437 to 504 of SEQ ID NO:4.
- 81. (New) The method of claim 68, wherein the TAB1 of (a) is a protein that binds to amino acids 76-303 of SEQ ID NO:2 and comprises an amino acid sequence encoded by a DNA that hybridizes with the complement of nucleotides 1338 to 1541 of SEQ ID NO:3 under washing conditions of 42°C, 5 x SSC, 0.1% sodium dodecyl sulfate, and 50% formamide.
- 82. (New) The method of claim 68, wherein the TAB1 of (a) is a protein that binds to amino acids 76-303 of SEQ ID NO:2 and comprises an amino acid sequence encoded by a DNA that hybridizes with the complement of nucleotides 1338 to 1541 of SEQ ID NO:3 under washing conditions of 60°C, 0.1 x SSC, and 0.1% sodium dodecyl sulfate.

Serial No.: 09/830,144 Filed: April 20, 2001

Page : 6 of 14

83. (New) The method of claim 68, wherein the inflammatory cytokine is IL-1, TNF, IL-10, or IL-6.

- 84. (New) The method of claim 68, wherein the inflammatory cytokine is IL-1.
- 85. (New) The method of claim 68, wherein the inflammatory cytokine is TNF.
- 86. (New) The method of claim 68, wherein the inflammatory cytokine is IL-6.
- 87. (New) The method of claim 68, wherein the inflammatory cytokine is IL-10.